

## Harnwegsinfekt Referenzen



### 11. Literatur

1. Walker M, Heady JA, Shaper AG: The prevalence of dysuria in woman in London. *J R Coll Gen Pract* 1983; 33: 411 - 415
2. Jolleys JV: Factors associated with regular episodes of dysuria among women in one rural general practice. *Br J Gen Pract* 1991; 41: 241-243
3. Diagnosis and Therapy survey 1996, Apoteksbolaget, Stockholm, 1997
4. Rasanathan M: Dysuria in general practice. *N Z Med J* 1979; 89: 54-56
5. Wathne B, Hovelius B, Mardh PA: Causes of frequency and dysuria in women. *Scand J Infect Dis* 1987; 19: 223-229
6. Jellheden B, Norrby R, Sandberg T: Symptomatic urinary tract infection in women in primary health care. *Scand J Prim Health Care* 1996; 14: 122-128
7. Barnett BJ, Stephens, DS: Urinary tract infection: an overview. *Am J Med Sci* 1997; 314: 245-249
8. Hooton TM, Scholes D, Hughes JP, et al.: A prospective study of risk factors for symptomatic urinary tract infection in young women. *N Engl J Med* 1996; 335: 468-474
9. Sheinfeld J, Schaeffer AJ, Cordon-Cardo C, et al.: Association of Lewis bloodgroup phenotye with recurrent urinary tract infection in women. *N Engl J Med* 1989; 320: 773-777
10. Smith HS, Hughes JP, Hooton TM, et al.: Antecedent antimicrobial use increases the risk of uncomplicated cystitis in young women. *Clin Infect Dis* 1997; 25: 63-68
11. Ikaheime R, Siitonen A, Heiskanen T, et al.: Recurrence of urinary tract infection in a primary care setting: analysis of a 1-year follow-up of 179 women. *Clin Infect Dis* 1996; 22: 91-99
12. Leiner S: Recurrent urinary tract infections in otherwise healthy adult women. Rational strategies for work-up and management. *Nurse Pract* 1995; 20: 48
13. Syme RR: Urinary tract infection in adult women: morbidity without mortality. *Aust Fam Physician* 1976; 5: 335-339
14. Mackenzie JR, Murphy AV, Beattie TJ, Azmy AF: Guidelines for the management of acute urinary tract infection in childhood. *J R Coll Physicians* 1991; 25: 263
15. Ahmed SM, Swedlund SK: Evaluation and treatment of urinary tract infections in children. *Am Fam Physician* 1998 Apr 1;57(7):1573-80, 1583-4
16. n.n.: Practice parameter: the diagnosis, treatment, and evaluation of the initial urinary tract infection in febrile infants and young children. American Academy of Pediatrics. Committee on Quality Improvement. Subcommittee on Urinary Tract Infection. *Pediatrics* 1999; 103: 843-52
17. Grun L, Tassano-Smith J, Carder C, et al.: Comparison of two methods of screening for genital chlamydial infection in women attending general practice: cross sectional survey. *BMJ* 1997; 315: 226-230
18. Nelson ME: Prevalence of Chlamydia trachomatis infection among women in a multiphysician primary care practice. *Am J Prev Med* 1992; 8: 298-302
19. Stamm WE, Hooton TM: Management of urinary tract infection in adults. *N Engl J Med* 1993; 329: 1328-1334
20. Huppertz BJ: Psychosomatic aspects of irritable bladder. *Urologe* 1986; 25: 84-49
21. Baldoni F, Baldaro B, Ercolani M, et al.: Urethral syndrome: a study in psychosomatic urology. *Psychother Psychosom* 1989; 52: 114-118
22. Chertok L, Bourguignon O, Guillon F, et al.: Urethral syndrome in the female ("irritable bladder"): the expression of fantasies about the urogenital area. *Psychosom Med* 1977; 39: 1-10
23. Sumners D, Kelsey M, Chait I: Psychological aspects of lower urinary tract infections in women. *BMJ* 1992; 304: 17-19
24. Nazareth I, King MB: The urethral syndrome: a controlled evaluation. *J Psychosom Res* 1993; 37: 737-743
25. Tait J, Peddie BA, Bailey RR, et al.: Urethral syndrome (abacterial cystitis) -search for a pathogen. *Br J Urol* 1985; 57: 552-556
26. Stamm WE, Wagner KF, Amsel R, et al.: Causes of the acute urethral syndrome in women. *N Engl J Med* 1980; 303: 409-415
27. Wilkins EG, Payne SR, Pead PJ, et al.: Interstitial cystitis and the urethral syndrome: a possible answer. *Br J Urol* 1989; 64: 39-44
28. Keay S, Schwalbe RS, Trifillis AL, et al.: A prospective study of microorganisms in urine and bladder biopsies from interstitial cystitis patients and controls. *Urology* 1995; 45: 223-229
29. Gittes RF, Nakamura RM: Female urethral syndrome. A female prostatitis? *West J Med* 1996; 164: 435-438
30. Bernstein AM, Philips HC, Linden W, et al.: A psychophysiological evaluation of female urethral

- syndrome: evidence for a muscular abnormality. *J Behav Med* 1992; 15: 299-312
31. Merkle W: Entzündungen. in: *Urologie*, Hrsg W. Merkle, Hippokrates Verl. Stuttgart, 1997, S. 116-147
  32. Rink E: Risk factors for urinary tract symptoms in women: beliefs among general practitioners and women and the effect on patient management. *Br J Gen Pract* 1998; 48: 1155-1158
  33. Bailey BL: Urinalysis predictive of urine culture results. *J Fam Pract* 1995; 40: 45-50
  34. Jellheden B, Norrby R, Sandberg T: Symptomatic urinary tract infection in women in primary health care. *Scand J Prim Health Care* 1996; 14: 122-128
  35. Hiraoka M, Hida Y, Hori C, Tuchida S, Kuroda M, Sudo M: Rapid dipstick test for diagnosis of urinary tract infection. *Acta Paediatr Jpn* 1994; 36: 379-382
  36. Baerheim A, Laerum E: Home-voided urine specimen in women. Diagnostic agreement with clean-catch midstream specimens. *Scand J Prim Health Care* 1990; 8: 207-211
  37. Immergut MA, Gilbert EC, Frensilli FJ, Goble M: The myth of the clean catch urine specimen. *Urology* 1981; 17: 339-340
  38. Lipsky BA, Inui TS, Plorde JJ, Berger RE: Is the clean-catch midstream void procedure necessary for obtaining urine culture specimens from men? *Am J Med* 1984; 76: 257-262
  39. Baerheim A, Digranes A, Hunskaar S: Evaluation of urine sampling technique: bacterial contamination of samples from women students. *Br J Gen Pract* 1992; 42: 241-243
  40. Goldsmith BM, Campos JM: Comparison of urine dipstick, microscopy, and culture for the detection of bacteriuria in children. *Clin Pediatr* 1990; 29: 214-8
  41. Winkens RA, Leffers P, Trienekens TA, Stobberingh EE: The validity of urine examination for urinary tract infections in daily practice. *Fam Pract* 1995; 12: 290-3
  42. Hiraoka M, Hida Y, Hori C, Tsuchida S, Kuroda M, Sudo M: Urine microscopy on a counting chamber for diagnosis of urinary infection. *Acta Paediatr Jpn* 1995; 37: 27-30
  43. Vickers D, Ahmad T, Coulthard MG: Diagnosis of urinary tract infection in children: fresh urine microscopy or culture? *Lancet* 1991; 338: 767-70
  44. Christiaens TCM, Meyere MD, Derese A: Disappointing specificity of the leucocyte-esterase test for the diagnosis of urinary tract infection in general practice. *Eur J Gen Pract* 1998; 4: 144-147
  45. Schaller P, Raetzo MA, Pittet D: Ich habe Brennen beim Wasserlassen. in: *Alltagsbeschwerden*, Hrsg. M.-A. Raetzo, A. Restellini, Verlag Hans Huber, Bern, 1998, S. 385-394
  46. Arav-Boger R, Leibovici L, Danon YL: Urinary tract infections with low and high colony counts in young women. Spontaneous remission and single-dose vs multiple-day treatment. *Arch Intern Med* 1994; 154: 300-304
  47. Kunin CM, White LV, Hua TH: A reassessment of the importance of "low-count" bacteriuria in young women with acute urinary symptoms. *Ann Intern Med* 1993; 119: 454-460
  48. Fowler JE, Pulaski ET: Excretory urography, cystography and cystoscopy in the evaluation of women with urinary-tract infection. *N Engl J Med* 1981; 304: 462-465
  49. Lipsky B: Prostatitis and urinary tract infection in men: what's new; what's true? *Am J Med* 1999; 106: 327-334
  50. Furrer HJ, Malinverni R: Harnwegsinfektionen bei Erwachsenen: alte und neue Aspekte. *Ther Umschau* 1994; 52: 842-852
  51. Pfau A, Sacks T, Engelstein D: Recurrent urinary tract infections in premenopausal women: prophylaxis based on an understanding of the pathogenesis. *J Urol* 1983 Jun; 129(6): 1153-7
  52. Baerheim A, Laerum E: Symptomatic lower urinary tract infection induced by cooling of the feet. *Scand J Prim Health Care* 1992; 10: 157-160
  53. Aune A, Alraek T, LiHua H, et al.: Acupuncture in the prophylaxis of recurrent lower urinary tract infection in adult women. *Scand J Prim Health Care* 1998; 16: 37-39
  54. Arzneimittelkommission der deutschen Ärzteschaft (Herausgeber): *Harnwegsinfektionen*. in: *Arzneiverordnungen*, 18. Auflage, Dt. Ärzte-Verlag Köln, 1997; 558-560
  55. British Medical Association / Royal Pharmaceutical Society of Great Britain: *British National Formulary No 35*. The Pharmaceutical Press, Oxon 1998
  56. Uhari M, Nuutinen M, Turtinen J: Adverse reactions in children during long-term antimicrobial therapy. *Pediatr Infect Dis J* 1996; 15: 404-408
  57. Brumfitt W, Hamilton-Miller JMT: Efficacy and safety profile of long-term nitrofurantoin in urinary tract infections: 18 years' experience. *J Antimicrob Chemother* 1998; 42: 363-371
  58. Irvani A, Klimberg I, Briefer C, et al.: A trial comparing low-dose, short-course ciprofloxacin and standard 7 day therapy with co-trimoxazole or nitrofurantoin in the treatment of uncomplicated urinary tract infection. *J Antimicrob Chemother* 1999; 43 Suppl A: 67-75
  59. Reeves DS: A perspective on the safety of antibacterials used to treat urinary tract infections. *J Antimicrob Chemother* 1994; 33 Suppl A: 111-120
  60. Christiaens TH, Heytens S, Verschraegen G, De Meyere M, De Maeseneer J: Which bacteria are found in Belgian women with uncomplicated urinary tract infections in primary health care, and what is their susceptibility pattern anno 95-96? *Acta Clin Belg* 1998; 53: 184-8
  61. van der Does MC, van Duijn NP, Timmerman CP, Degener JE: Resistance to antibiotics in uncomplicated urinary tract infections. *Huisarts Wet* 1998; 41: 421-3
  62. Preston St, Abdel-Rahman SM, Nahata MC: Empiric treatment of uncomplicated urinary tract infections. *Ann Pharmacother* 1998; 32: 1231-1233
  63. Gupta K, Scholes D, Stamm WE: Increasing prevalence of antimicrobial resistance among uropathogens causing uncomplicated cystitis in women. *JAMA* 1999; 281: 736-738
  64. Fernandez Fernandez A, Lantero Benedito M, Gastanares Hernando MJ, Undabeitia Santisteban E,

- Fernandez Santos F, Castano Rodriguez A, et al.: Quinolone-resistant *Escherichia coli* in the health area of a 650-bed hospital. *Actas Urol Esp* 1994; 18: 634-8
65. Osterlund A, Olsson-Liljequist B: Fluoroquinolone resistance of human pathogenic bacteria. Resistant *E coli* now appearing in Sweden. *Lakartidningen* 1999 Apr 21; 96(16): 1965-6
  66. Webster CA, Curran R, Towner KJ: Comparative in -vitro activity of cefaclor against urinary tract isolates of *Escherichia coli*. *J Antimicrob Chemother* 1996; 38: 59-66
  67. Stamm WE: Controversies in single dose therapy of acute uncomplicated urinary tract infections in women. *Infection* 1992; 20: 272-275
  68. Raz R, Rottensterich E, Leshem Y, et al.: Double-blind study comparing 3-day regimens of cefixime and ofloxacin in treatment of uncomplicated urinary tract infections in women. *Antimicrob Agents Chemother* 1994; 38: 176-1177
  69. Keenan TD, Elliott JC, Bishop V, Peddie BA, Bailey RR: Comparison of trimethoprim alone with co-trimoxazole and sulphamethizole for treatment of urinary tract infections. *N Z Med J* 1983; 96: 341-342
  70. Ziak E, Sedlmayr P, Sterz F: Trimethoprim as a monosubstance and cotrimoxazole in infections of the efferent urinary tract. *Wien Med Wochenschr* 1987; 137: 123-126
  71. Trimethoprim Study Group: Comparison of trimethoprim at three dosage levels with co-trimoxazole in the treatment of acute symptomatic urinary tract infection in general practice. *J Antimicrob Chemother* 1981; 7: 179-183
  72. Lacey RW, Lord VL, Gunasekera HK, et al.: Comparison of trimethoprim alone with trimethoprim sulphamethoxazole in the treatment of respiratory and urinary infections with particular reference to selection of trimethoprim resistance. *Lancet* 1980; 1: 1270-1273
  73. Jick H: Adverse reactions to trimethoprim-sulphamethoxazole in hospitalized patients. *Rev Infect Dis* 1982; 4: 426-428
  74. Hunziker T, Kunzi UP, Braunschweig S, Zehnder R, Hoigne R: Comprehensive hospital drug monitoring (CHDM): adverse skin reactions, a 20-year survey. *Allergy* 1997; 52: 388-393
  75. Lee AJ, Maddix DS: Trimethoprim/sulfamethoxazole-induced hypoglycemia in a patient with acute renal failure. *Ann Pharmacother* 1997; 31: 727-732
  76. Schopf E: Skin reactions to co-trimoxazole. *Infection* 1987; 15: S254-S258
  77. Anonymous: Cotrimoxazole use restricted. *Drug Therap Bull* 1995; 33: 92-93
  78. Ponte CD, Fisher MA: Use of fluoroquinolones: practical considerations. *Am Fam Physician* 1993; 47: 1243-1249
  79. Stratton C: Fluoroquinolone antibiotics: properties of the class and individual agents. *Clin Ther* 1992; 14: 348-375
  80. Van Pienbroek E, Hermans J, Kaptein AA, Mulder JD: Fosfomycin trometamol in a single dose versus seven days nitrofurantoin in the treatment of acute uncomplicated urinary tract infections in women. *Pharm World Sci* 1993; 15: 257-62
  81. Stein GE: Single-dose treatment of acute cystitis with fosfomycin trometamol. *Ann Pharmacother* 1998; 32: 215-9
  82. Avorn J, Monane M, Gurwitz JH, et al.: Reduction of bacteriuria and pyuria after ingestion of cranberry juice. *JAMA* 1994; 271: 751-754
  83. Walker EB, Barney DP, Mickelsen JN, et al.: Cranberry concentrate: UTI prophylaxis. *J Fam Pract* 1997; 45: 167-168
  84. Jepson RG, Mihaljevic L, Craig J: Cranberries for preventing urinary tract infections (Cochrane Review). *Cochrane Library* 1999; 1
  85. Schulman CC, Corbusier A, Michiels H, Taenzer HJ: Oral immunotherapy of recurrent urinary tract infections: a double-blind placebo-controlled multicenter study. *J Urol* 1993; 150: 917-921
  86. Tammen H, German Urinary Tract Infection Study Group: Immunobiotherapy with Uro-Vaxom in recurrent urinary tract infection. *Br J Urol* 1990; 65: 6-9
  87. Hooton TM, Wintwer C, Tiu F, et al.: Randomized comparative trial and cost analysis of 3-day antimicrobial regimens for treatment of acute cystitis in women. *JAMA* 1995; 273: 41-45
  88. Gossius G, Vorland L: A randomized comparison of single-dose vs three-day and ten-day therapy with trimethoprim-sulfamethoxazole for acute cystitis in women. *Scand J Infect Dis* 1984; 16: 373-379
  89. Norrby SR: Short-term treatment of uncomplicated lower urinary tract infections in women. *Rev Infect Dis* 1990; 12: 458-467
  90. Österberg, Aberg H, Hallander HO, et al.: Efficacy of single-dose versus seven-day trimethoprim treatment of cystitis in women. *J Inf Dis* 1990; 161: 942-947
  91. Greenberg RN, Reilly PM, Luppen KL, et al.: Randomized study of single-dose, three-day, and seven day treatment of cystitis in women. *J Infect Dis* 1986; 153: 277-282
  92. Fihn SD, Johnson C, Roberts PL et al.: Trimethoprim-sulphamethoxazole for acute dysuria in women: a single-dose or 10-day course. A double-blind, randomized trial. *Ann Int Med* 1988; 108: 350-357
  93. Nicolle LE: Urinary tract infection in the elderly - how to treat and when. *Infection* 1992; 20: 261-265
  94. Schaeffer AJ, Stuppy BA: Efficacy and safety of self-start therapy in women with recurrent urinary tract infections. *J Urol* 1999 Jan;161(1): 207-11
  95. Stapleton A, Latham RH, Johnson C, et al.: Postcoital antimicrobial prophylaxis for recurrent urinary tract infection. A randomized, double-blind, placebo-controlled trial. *JAMA* 1990; 264: 703-706
  96. Stamm WE, Counts GW, Wagner KF, et al.: Antimicrobial prophylaxis of recurrent urinary tract infections: a double-blind, placebo-controlled trial. *Ann Intern Med* 1980; 92: 770-775
  97. Nicolle LE, Harding GK, Thomson M, et al.: Efficacy of five years of continuous, low-dose trimethoprim-sulfamethoxazole prophylaxis for urinary tract infection. *J Infect Dis* 1988; 157: 1239-

- 1242
98. Raz R, Stamm WE: A controlled trial of intravaginal estriol in postmenopausal women with recurrent urinary tract infections. *N Engl J Med* 1993; 329: 753-6
  99. Vercaugne LM, Zhanel G: Recommended treatment for urinary tract infection in pregnancy. *Ann Pharmacother* 1994; 28: 248-251
  100. Smail F: Antibiotic vs no treatment for asymptomatic bacteriuria in pregnancy. *Cochrane Library* 1996
  101. DiGiuseppe C, United States Preventive Services Task Force: Guide to clinical preventive services. 2nd ed. Baltimore (MD): Williams & Wilkins; 1996; 347-59
  102. Warren JW, Anthony WC, Hoopes JM, et al.: Cephalexin for susceptible bacteriuria in afebrile, long-term catheterized patients. *JAMA* 1982; 248: 454-458
  103. Ouslander JG, Schapira M, Schnelle JF, et al.: Does eradicating bacteriuria affect the severity of chronic urinary incontinence in nursing home residents. *Ann Int Med* 1995; 122: 749-754
  104. Abrutyn E, Mossey J, Berlin J, et al.: Does asymptomatic bacteriuria predict mortality and does antimicrobial treatment reduce mortality in elderly ambulatory women? *Ann Int Med* 1994; 120: 827-833
  105. Ronald AR, Nicolle LE, Harding GKM: Standards of therapy for urinary tract infections in adults. *Infection* 1992; 20: 164-170
  106. Grun L, Tassano-Smith J, Carder C, Johnson AM, Robinson A, Murray E, et al.: Comparison of two methods of screening for genital chlamydial infection in women attending in general practice: cross sectional survey. *BMJ* 1997; 315: 226-230
  107. Lauharanta J, Saarinen K, Mustonen MT, Happonen HP.: Single-dose azithromycin versus seven -day doxycyclin in the treatment of non-gonococcal urethritis in males. *J Antimicrob Chemother* 1993; 31 (Suppl. E): 177-83
  108. Nilsen A, Halsos A, Johansen A, et al.: A double blind study of single doses azythromycin and doxycyclin in the treatment of chlamydial urethritis in males. *Genitourin Med* 1992; 68: 325-27
  109. Martin DH, Mroczkowski TF, Dalu ZA, et al.: A controlled trial of a sinle dose of azithromycin for the treatment of urethritis and cervicitis. The Azithromycin for Chlamydial Inections Study Group. *N Engl J Med* 1992; 327: 921-25
  110. Adair CD, Gunter M, Stovall TG, McElroy G, Veille JC, Ernest JM: Chlamydia in pregnancy: a randomized trial of azithromycin and erythromycin. *Obstet Gynecol* 1998; 91: 165-68
  111. Stenberg A, Heimer G, Ulmsten U: The prevalence of urogenital symptoms in postmenopausal women. *Maturitas* 1995; 22: S17-S20
  112. Günther EA: Psychosomatische Urologie. in: Urologie, Hrsg W. Merkle, Hippokrates Verl. Stuttgart, 1997, S. 393-404
  113. Parzani S, Constantini E, Petroni PA, et al.: Urethral syndrome: clinical results with antibiotics alone or combined with estrogen. *Eur Urol* 1994; 26: 115-119
  114. Oxman AD: Checklists for review articles. *BMJ* 1994; 309: 648-651
  115. Barry HC, Ebell MH, Hickner J: Evaluation of suspected urinary tract infection in ambulatory women: A cost-utility analysis of office-based strategies. *J Fam Pract* 1997; 44: 49-60
  116. Barwitz HJK: Leitsymptome Dysurie. Versuch einer Rationalisierung des Vorgehens in der Allgemeinpraxis mit einer Handlungsleitlinie. *Münch med Wochenschrift* 1999; 141(12): 147-150

© Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin  
AWMF-Register

Link 1 CochraneReview über Cranberries zur Prophylaxe <http://www.update-software.com/ccweb/cochrane/revabstr/ab001321.htm>

zur Therapie: <http://www.update-software.com/ccweb/cochrane/revabstr/ab001322.htm>

[\[Start Leitlinien\]](#) [\[ » Asthma\]](#) [\[ » Kopfschmerzen\]](#) [\[ » Hypertonie\]](#) [\[ » Otitis media\]](#) [\[ » Gallensteine\]](#) [\[ » Herzinsuffizienz\]](#)  
[\[ » Demenz\]](#) [\[ » Harnwegsinfekt\]](#) [\[Algorithmus\]](#) [\[Volltext\]](#) [\[Referenzen\]](#) [\[Patientenleitlinie\]](#) [\[Impressum/ ©\]](#) [\[ » KR-](#)  
[Karzinom\]](#) [\[ » Dekubitus\]](#) [\[Ihre Meinung?\]](#) [\[Interne Infos\]](#)

Schon bekannt? Testen Sie Ihr Wissen und sammeln Sie CME-Punkte mit [Medizinerwissen.de](#)

© Medizinisches Wissensnetzwerk [evidence.de](#) der Universität Witten/Herdecke [Patienten...](#)

Update:17/06/05  [ju-ko.de](#)